

## Guidelines for the Blood Transfusion Services

### 7.4.1: Platelets, Pooled, Buffy Coat Derived, Leucocyte Depleted

<http://www.transfusionguidelines.org/red-book/chapter-7/7-4/7-4-1>

### 7.4.1: Platelets, Pooled, Buffy Coat Derived, Leucocyte Depleted

A pool of platelets, derived from buffy coats, which contains less than  $1 \times 10^6$  leucocytes.

#### 7.4.1.1: Technical information

- Donations of whole blood where the bleed time exceeded 15 minutes are not suitable for platelet production.
- The buffy coats must be prepared at ambient temperature from whole blood where the surface temperature of packs has not dropped below 18°C.
- Initial separation of buffy coat must occur within 24 hours of venepuncture (unless supported by additional validation), with a minimum buffy coat rest period of 2 hours before secondary pooling and processing of buffy coats to produce the final component, which is generally completed before the end of Day 1.
- The volume of suspension medium must be sufficient to maintain the pH at  $\geq 6.4$  at the end of the shelf life of the component.
- The production process transfers the final component into a pack that was not part of the original pack assembly. Therefore a secure system must be in place to ensure a full audit trail and that the correct identification number is put on the final component pack.
- Where the production method requires the use of a single unit of plasma for resuspension, the plasma from group O donors should be tested for high-titre anti-A and anti-B and 'high-titre negative' units labelled. The testing method and acceptable limits should be defined (see also Chapter 9). Plasma should be selected from male donors as a TRALI risk reduction strategy.
- Platelets, Pooled, Buffy Coat Derived, Leucocyte Depleted, should be administered through a CE /UKCA/UKNI marked transfusion set.

#### 7.4.1.2: Labelling

For general guidelines, see section 6.6.

The following shall be included on the label:

(\* = in eye-readable and UKBTS approved barcode format)

- Platelets, Pooled, Buffy Coat Derived, Leucocyte Depleted\* and volume
- the blood component producer's name\*

- a unique pool or batch number or the donation number of all contributing platelet units\*
- the ABO group\*
- the RhD group stated as positive or negative\*
- the expiry date\*
- the temperature of storage and a comment that continuous gentle agitation throughout storage is recommended
- the blood pack lot number\*
- the name, composition and volume of the anticoagulant or additive solution.

In addition, the following statements should be made:

#### **INSTRUCTION**

*Always check patient/component compatibility/identity*

*Inspect pack and contents for signs of deterioration or damage*

*Risk of adverse reaction/infection, including vCJD*

#### **7.4.1.3: Storage**

---

For general guidelines, see section 6.7.

- The storage period depends on a number of factors including the nature of the container, the concentration of platelets and on whether an open or closed system is used.
- Packs currently in use for this purpose allow for storage at a core temperature of  $22 \pm 2^\circ\text{C}$  with continuous gentle agitation for up to 5 days in a closed system. Appropriate pack and platelet concentration combinations may allow storage up to 7 days, but due to concerns over bacterial contamination requires either an assay to exclude bacterial contamination prior to transfusion or application of a licensed pathogen inactivation procedure.
- If any production stage involves an open system, after preparation the component should be used as soon as possible. If storage is unavoidable, the component should be stored at a core temperature of  $22 \pm 2^\circ\text{C}$  with continuous agitation and used within 6 hours.
- Platelets should be gently agitated during storage. If agitation is interrupted, for example due to equipment failure or prolonged transportation, the components are suitable for use, retaining the same shelf life, provided that no single interruption lasts for more than eight hours, and the total length of all interruptions is no longer than 24 hours.

#### **7.4.1.4: Testing**

---

In addition to the mandatory and other tests required for blood donations described in Chapter 9, and leucocyte counting (see sections 6.3 and 7.1.1), a minimum of 75% of those components tested for the parameters shown in Table 7.4.1 shall meet the specified values.

**Table 7.4.1 Platelets, Pooled, Buffy Coat Derived, Leucocyte Depleted – additional tests**

Parameter	Frequency of test	Specification
Volume <sup>1</sup>	1% or as determined by statistical process control (if $\leq 10$ components produced per month then test every available component)	Within locally defined nominal volume range
Platelet count <sup>2</sup>		$\geq 240 \times 10^9/\text{pool}$
pH at end of shelf life <sup>3</sup>		$\geq 6.4$
Leucocyte count <sup>4</sup>	As per sections 6.3 and 7.1.1	$< 1 \times 10^6/\text{pool}$
<sup>1</sup> Units measured and found to be $< 150$ mL or $> 380$ mL should only be issued for transfusion under concessionary release		
<sup>2</sup> Units measured and found to have $< 160 \times 10^9/\text{pool}$ , or more than the maximum recommended by the manufacturer of the storage pack where stated, should only be issued for transfusion under concessionary release		
<sup>3</sup> A minimum of 95% of those components tested shall meet the specified values		
<sup>4</sup> Methods validated for counting low numbers of leucocytes must be used		

*Note: Visual inspection of platelet components for the swirling phenomenon, clumping, excessive red cell contamination and abnormal volume is a useful pre-issue check.*

#### 7.4.1.5: Transportation

For general guidelines, see section 6.11.

- Containers for transporting platelets should be equilibrated at room temperature before use. During transportation the temperature of platelets must be kept as close as possible to the recommended storage temperature and, on receipt, unless intended for immediate therapeutic use, the component should be transferred to storage at a core temperature of  $22 \pm 2^\circ\text{C}$  with continuous gentle agitation.
- Plastic overwraps should be removed prior to storage.